

On the Algebraic Structure of Second Order Symmetric Tensors in 5-Dimensional Space-times

G.S. Hall*, M.J. Rebouças[†], J. Santos[‡] and A.F.F. Teixeira[§]

* Department of Mathematical Sciences
University of Aberdeen
Aberdeen AB9 2TY, Scotland – U.K.

^{† §} Centro Brasileiro de Pesquisas Físicas
Departamento de Relatividade e Partículas
Rua Dr. Xavier Sigaud 150
22290-180 Rio de Janeiro – RJ, Brazil

[‡] Universidade Federal do Rio G. do Norte
Departamento de Física, Caixa Postal 1641
59072-970 Natal – RN, Brazil

February 7, 2008

Abstract

A new approach to the algebraic classification of second order symmetric tensors in 5-dimensional space-times is presented. The possible Segre types for a symmetric two-tensor are found. A set of canonical forms for each Segre type is obtained. A theorem which collects together some basic results on the algebraic structure of the Ricci tensor in 5-dimensional space-times is also stated.

*INTERNET: GSH@MATHS.ABDN.AC.UK

[†]INTERNET: REBOUCAS@CAT.CBPF.BR

[‡]INTERNET: JANILO@DFTE.UFRN.BR

[§]INTERNET: TEIXEIRA@CAT.CBPF.BR

1 Introduction

In general relativity, it is well known that the curvature tensor can be uniquely decomposed into three irreducible parts, namely the Weyl tensor, the traceless Ricci tensor and the Ricci scalar. Petrov and others [1] – [4] have discussed the algebraic classification of the Weyl part of the curvature tensor. The Petrov classification as it is known nowadays has played an important role in the investigation of various issues in general relativity [5].

The algebraic classification of the Ricci part, known as the Segre classification, has been discussed by several authors [6] and is of great importance in three contexts. One is in understanding some purely geometrical features of space-times [7] – [10]. The second one is in classifying and interpreting matter field distributions [11] – [16]. The third is as part of the procedure for checking whether apparently different space-times are in fact locally the same up to coordinate transformations (the equivalence problem [17] – [22]).

In a recent article Santos *et al.* [23] have studied the algebraic classification of second order symmetric tensors defined on 5-dimensional (5-D for short) Lorentz manifolds M , extending previous results for 3-D and 4-D space-times [24] – [26]. Their analysis is made from first principles, i.e., without using the previous classifications on lower dimensional space-times. However, as concerns the classification itself their approach is not straightforward.

In this work we examine the algebraic structure of second order symmetric tensors defined on 5-dimensional space-times under different bases. We shall assume the algebraic classification of the Ricci tensor for 4-D space-times, and then show that the algebraic classification Santos *et al.* [23] have studied can be achieved in a considerably simpler way. Their results are, therefore, recovered under different premises. We also state a fair number of new results concerning the algebraic structure of the Ricci tensor and collect them together in a theorem, which generalizes theorems on 4-D space-times [6, 9, 24].

2 Prerequisites

In this section we briefly review some important results required for the remainder of this work. For a detailed account of the notation used throughout this article see [6, 23].

The classification of a second order symmetric real tensor at points on a 5-dimensional manifold carrying a Lorentz metric of signature $(-++++)$ can be obtained directly from the analogous classification in the 4-dimensional case where the results are known [2, 6, 8, 24]. The situation in the latter case can now be summarized in the following theorem in which $\mathbf{t}, \mathbf{x}, \mathbf{y}, \mathbf{z}$ and $\mathbf{l}, \mathbf{m}, \mathbf{x}, \mathbf{y}$ denote, respectively, a pseudo-orthonormal and a null tetrad and in which the only non-vanishing inner products are $-t_a t^a = x_a x^a = y_a y^a = z_a z^a = 1$ and $l_a m^a = x_a x^a = y_a y^a = 1$, respectively.

Theorem 2.1 *Let (M, g) be a space-time so that M is a 4-dimensional manifold and the metric g has Lorentz signature $(-+++)$. Let $p \in M$ and let T be a real symmetric second order tensor at p . Then (with respect to g) T must take one of the Segre types $\{1, 111\}$, $\{211\}$, $\{31\}$ and $\{z \bar{z} 11\}$ or a degeneracy thereof. The corresponding canonical forms in some appropriate tetrad are:*

(i) *for the type $\{1, 111\}$ either of*

$$T_{ab} = 2\alpha_1 l_{(a} m_{b)} + \alpha_2 (l_a l_b + m_a m_b) + \alpha_3 x_a x_b + \alpha_4 y_a y_b, \quad (2.1)$$

$$T_{ab} = (\alpha_2 - \alpha_1) t_a t_b + (\alpha_1 + \alpha_2) z_a z_b + \alpha_3 x_a x_b + \alpha_4 y_a y_b, \quad (2.2)$$

(ii) *for the type $\{211\}$*

$$T_{ab} = 2\beta_1 l_{(a} m_{b)} \pm l_a l_b + \beta_2 x_a x_b + \beta_3 y_a y_b, \quad (2.3)$$

(iii) *for the type $\{31\}$*

$$T_{ab} = 2\gamma_1 l_{(a} m_{b)} + 2l_{(a} x_{b)} + \gamma_1 x_a x_b + \gamma_2 y_a y_b, \quad (2.4)$$

(iv) *finally for the type $\{z \bar{z} 11\}$*

$$T_{ab} = 2\delta_1 l_{(a} m_{b)} + \delta_2 (l_a l_b - m_a m_b) + \delta_3 x_a x_b + \delta_4 y_a y_b, \quad (2.5)$$

where $\alpha_1, \alpha_2, \dots, \delta_4 \in \mathbb{R}$ and $\delta_2 \neq 0$.

3 The Algebraic Classification

Now suppose M is a 5-dimensional manifold carrying a Lorentz metric g and, with $p \in M$, let T be a real second order, symmetric tensor at p with components T^{ab} in some basis of the tangent space $T_p M$ to M at p . First note that any non-trivial proper subspace of $T_p M$ may be classified into one of the three types, *spacelike*, *null* and *timelike* according as it contains no null directions, a unique null direction or more than one null direction, respectively. Further, the (g) -orthogonal complement of a spacelike (timelike) subspace is timelike (spacelike). Next, if the tensor T , when regarded as a linear transformation from $T_p M$ to itself with matrix T^a_b , admits a non-null invariant subspace V of $T_p M$ then the orthogonal complement of V is also an invariant subspace of T as is easily checked (see [6] for the case when (M, g) is a space-time). Now the tensor T *must* admit an invariant 2-dimensional subspace. To see this note that if all the eigenvalues of T are real then the first two members (in the conventional numbering) of a canonical Jordan basis span such a subspace whilst if T admits a complex eigenvalue the real and imaginary parts of any corresponding complex eigenvector span an invariant subspace for T .

Lemma 3.1 *In the above notation T must admit a spacelike eigenvector.*

Proof Suppose first that T has a complex eigenvalue $\lambda = a + ib$ ($a, b \in \mathbb{R}$, $b \neq 0$) and corresponding complex eigenvectors $\mathbf{k} = \mathbf{r} + i\mathbf{s}$ ($\mathbf{r}, \mathbf{s} \in T_p M$, $\mathbf{s} \neq 0$). Then

$$T_{ab}(r^b + is^b) = (a + ib)(r_a + is_a) . \quad (3.1)$$

By replacing $\mathbf{k}$ with $z\mathbf{k}$ ($z \in \mathbb{C}$), if necessary, one can arrange that $s_a r^a = 0$. On contracting the real part of (3.1) with s^a and the imaginary part with r^a one sees that $r_a r^a + s_a s^a = 0$. Now $\mathbf{r}$ and $\mathbf{s}$ cannot both be null for then they would be parallel and (3.1) would give $b = 0$. Hence one of $\mathbf{r}$ and $\mathbf{s}$ is timelike and the other spacelike and so the invariant subspace they span is timelike. The orthogonal complement of this subspace is thus spacelike and the restriction of T to it is a standard self adjoint Euclidean action and

is diagonalisable. It follows that T has a spacelike eigenvector lying in this orthogonal complement.

Now suppose T has only real eigenvalues. If T admits a spacelike invariant 2-dimensional subspace the above argument applies, whilst if T admits a timelike invariant 2-dimensional subspace the above argument applies to its (invariant) orthogonal complement. So suppose that T admits only null invariant 2-dimensional subspaces. The unique null direction in each such null invariant subspace is, in fact, an eigendirection of T (see [6] for a proof in the 4-dimensional space-time case). Let $\mathbf{k}_1$ be such a null eigendirection of T with eigenvalue $\mu_1 \in \mathbb{R}$. If T admits any other (independent) eigenvector $\mathbf{k}_2$ with eigenvalue μ_2 then $\mathbf{k}_2$ and μ_2 are real since, by assumption, T admits only real eigenvalues. But then either $\mu_1 = \mu_2$ (in which case the subspace of $T_p M$ spanned by $\mathbf{k}_1$ and $\mathbf{k}_2$ consists entirely of eigenvectors and must contain a spacelike member) or else $\mu_1 \neq \mu_2$, in which case $\mathbf{k}_1$ and $\mathbf{k}_2$ are orthogonal (since T is symmetric) and so $\mathbf{k}_2$ is spacelike, since $\mathbf{k}_1$ is null.

Thus the only remaining possibility is when T has the single (independent) eigenvector $\mathbf{k}_1$ which is real and null. The Segre type for T is then $\{5\}$ and its Jordan form in some (Jordan) basis at p is

$$T^a_b = \begin{bmatrix} \mu_1 & 1 & 0 & 0 & 0 \\ 0 & \mu_1 & 1 & 0 & 0 \\ 0 & 0 & \mu_1 & 1 & 0 \\ 0 & 0 & 0 & \mu_1 & 1 \\ 0 & 0 & 0 & 0 & \mu_1 \end{bmatrix}.$$

The condition that $g_{ac}T^c_b$ is symmetric is then easily shown to imply that $g_{12} = g_{22} = 0$ at p . Thus the first two (column) vectors in the Jordan basis are null and orthogonal at p , contradicting the Lorentz signature of g , and so this case cannot occur. This completes the proof.

Theorem 3.1 *Let M be a 5-dimensional manifold admitting a Lorentz metric g with signature $(- + + + +)$. Let $p \in M$ and let T_{ab} be a real symmetric second order tensor at p . Then (with respect to g) T must take one of the Segre types $\{1, 1111\}$, $\{2111\}$, $\{311\}$ and $\{z \bar{z}111\}$ or a degeneracy thereof. The corresponding canonical forms in some appropriate pentad (either orthonormal $\mathbf{t}, \mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{w}$ or null $\mathbf{l}, \mathbf{m}, \mathbf{x}, \mathbf{y}, \mathbf{w}$ in which the only non-vanishing inner products are, respectively, $-t_a t^a = x_a x^a = y_a y^a = z_a z^a = w_a w^a = 1$ and $l_a m^a = x_a x^a = y_a y^a = w_a w^a = 1$) at p are:*

(i) *for the type $\{1, 1111\}$ either of*

$$T_{ab} = 2\alpha_1 l_{(a} m_{b)} + \alpha_2 (l_a l_b + m_a m_b) + \alpha_3 x_a x_b + \alpha_4 y_a y_b + \alpha_5 w_a w_b, \quad (3.2)$$

$$T_{ab} = (\alpha_2 - \alpha_1) t_a t_b + (\alpha_1 + \alpha_2) z_a z_b + \alpha_3 x_a x_b + \alpha_4 y_a y_b + \alpha_5 w_a w_b, \quad (3.3)$$

(ii) *for the type $\{2111\}$*

$$T_{ab} = 2\beta_1 l_{(a} m_{b)} \pm l_a l_b + \beta_2 x_a x_b + \beta_3 y_a y_b + \beta_4 w_a w_b, \quad (3.4)$$

(iii) *for the type $\{311\}$*

$$T_{ab} = 2\gamma_1 l_{(a} m_{b)} + 2l_{(a} x_{b)} + \gamma_1 x_a x_b + \gamma_2 y_a y_b + \gamma_3 w_a w_b, \quad (3.5)$$

(iv) *finally for the type $\{z \bar{z}111\}$*

$$T_{ab} = 2\delta_1 l_{(a} m_{b)} + \delta_2 (l_a l_b - m_a m_b) + \delta_3 x_a x_b + \delta_4 y_a y_b + \delta_5 w_a w_b, \quad (3.6)$$

where $\alpha_1, \alpha_2, \dots, \delta_5 \in \mathbb{R}$ and $\delta_2 \neq 0$.

Proof By the lemma T admits a spacelike eigenvector, say $\mathbf{w}$ at p scaled so that $w_a w^a = 1$. The orthogonal complement of $\mathbf{w}$ in $T_p M$ is a 4-dimensional subspace V of $T_p M$ which is timelike (i.e. it admits the restriction of $g(p)$ as a Lorentz metric) and invariant under T and so T has a self adjoint action on V . So the restriction of T to V is controlled by the previous theorem and, accordingly, some appropriate tetrad may be chosen in V to achieve one of the canonical forms (2.1) – (2.5) for this restriction. Appending the vector $\mathbf{w}$ to these tetrads yields pentads in which T takes the canonical forms (3.2) – (3.6).

4 Further Results and Concluding Remarks

In this section we shall present some results concerning the algebraic structure of a second order symmetric tensor T defined on a 5-dimensional Lorentz manifold M , which generalizes theorems on 4-dimensional space-times [6, 8, 9, 24] and which can be collected in the following theorem:

Theorem 4.1 *Let M be a real 5-dimensional manifold endowed with a Lorentz metric g of signature $(- + + + +)$. Let T be a real second order symmetric tensor defined at a point $p \in M$. Then*

- (i) *T has a timelike eigenvector if and only if it is diagonalizable over $\mathbb{R}$ (type $\{1, 1111\}$).*
- (ii) *T has at least three real orthogonal independent eigenvectors, two of which (at least) are spacelike.*
- (iii) *T has all eigenvalues real and is not diagonalizable if and only if it has an unique null eigendirection.*
- (iv) *If T has two linearly independent null eigenvectors then it is diagonalizable over $\mathbb{R}$ (type $\{1, 1111\}$).*
- (v) *There always exists a 2-D spacelike subspace of $T_p M$ invariant under T .*
- (vi) *If a non-null subspace of $T_p M$ is invariant under T , then so is its orthogonal complement.*
- (vii) *There always exists a 3-D timelike subspace of $T_p M$ invariant under T .*
- (viii) *If T admits a null invariant subspace $\mathcal{N}$ of any dimension n ($=2, 3$, or 4) then it admits one of each dimension 2, 3 and 4, and this occurs if and only if T admits a null eigenvector (which necessarily lies in any such $\mathcal{N}$ admitted).*

The proofs can be gathered essentially by inspection of the canonical forms (3.2) – (3.6) and are not presented here for the sake of brevity (for proofs of similar theorems in 4-dimensional space-times see [6]).

To conclude, we remark that by a similar procedure to that used in the lemma 3.1 one can show that a symmetric two-tensor T defined on an n -dimensional ($n \geq 4$) Lorentz space has at least one real *spacelike* eigenvector. The existence of this eigenvector can be

used to reduce, by induction, the classification of symmetric two-tensors on n -dimensional ($n \geq 4$) spaces to the classification on 4-dimensional spaces, thus recovering in a simpler way the results of [27].

References

- [1] A. Z. Petrov, *Sci. Notices of Kazan State University* **114**, 55 (1954).
- [2] A. Z. Petrov, “Einstein Spaces”, Pergamon Press (1969).
- [3] R. Penrose, *Ann. Phys. NY* **10**, 171 (1960).
- [4] R. Penrose and W. Rindler , “Spinors and Space-Time”, vol. 2, Cambridge U. P., Cambridge (1986).
- [5] See, for example, D. Kramer, H. Stephani, M. MacCallum and E. Herlt, “Exact Solutions of Einstein’s Equations”, Cambridge U. P., Cambridge (1980).
- [6] G. S. Hall, *Diff. Geom.* **12**, 53 (1984). This reference contains an extensive bibliography on the classification of the Ricci tensor on 4-dimensional space-times.
- [7] R. V. Churchill, *Trans. Amer. Math. Soc.* **34**, 784 (1932).
- [8] J. Plebański, *Acta Phys. Pol.* **26**, 963 (1964).
- [9] W. J. Cormack and G. S. Hall, *J. Phys. A* **12**, 55 (1979).
- [10] C. Bona, B. Coll and J. A. Morales, *J. Math. Phys.* **33**, 670 (1992).
- [11] G. S. Hall and D. A. Negm, *Int. J. Theor. Phys.* **25**, 405 (1986).
- [12] G. S. Hall, *Arab. J. Sci. Eng.* **9**, 87 (1984).
- [13] J. J. Ferrando, J.A. Morales and M. Portilla, *Gen. Rel. Grav.* **22**, 1021 (1990).

- [14] M. J. Rebouças and A. F. F. Teixeira, *J. Math. Phys.* **32**, 1861 (1991).
- [15] M. J. Rebouças and A. F. F. Teixeira, *J. Math. Phys.* **33**, 2855 (1992).
- [16] J. Santos, M. J. Rebouças and A. F. F. Teixeira, *J. Math. Phys.* **34**, 186 (1993).
- [17] E. Cartan, “Leçons sur la Géométrie des Éspaces de Riemann”, Gauthier-Villars, Paris (1951). Reprinted, Éditions Jacques Gabay, Paris (1988). English translation by J. Glazebrook, Math. Sci. Press, Brookline (1983).
- [18] A. Karlhede, *Gen. Rel. Grav.* **12**, 693 (1980).
- [19] See M. A. H. MacCallum and J. E. F. Skea, “SHEEP: A Computer Algebra System for General Relativity”, in *Algebraic Computing in General Relativity, Lecture Notes from the First Brazilian School on Computer Algebra*, Vol. II, edited by M. J. Rebouças and W. L. Roque. Oxford U. P., Oxford (1994); and references therein.
- [20] M. A. H. MacCallum, “Classifying Metrics in Theory and Practice”, in *Unified Field Theory in More Than 4 Dimensions, Including Exact Solutions*, edited by V. de Sabbata and E. Schmutzer. World Scientific Publishing Co., Singapore (1983).
- [21] M. A. H. MacCallum, “Algebraic Computing in General Relativity”, in *Classical General Relativity*, edited by W. B. Bonnor, J. N. Islam and M. A. H. MacCallum. Cambridge U. P., Cambridge (1984).
- [22] M. A. H. MacCallum, “Computer-aided Classification of Exact Solutions in General Relativity”, in *General Relativity and Gravitational Physics (9th Italian Conference)*, edited by R. Cianci, R. de Ritis, M. Francaviglia, G. Marmo, C. Rubano and P. Scudellaro. World Scientific Publishing Co., Singapore (1991).
- [23] J. Santos, M. J. Rebouças and A. F. F. Teixeira, *J. Math. Phys.* **36**, 3074 (1995).
- [24] G. S. Hall, *J. Phys. A* **9**, 541 (1976).

- [25] G. S. Hall, T. Morgan and Z. Perjés, *Gen. Rel. Grav.* **19**, 1137 (1987).
- [26] G. S. Hall, “Physical and Geometrical Classification in General Relativity”, Brazilian Centre for Physics Research Monograph, CBPF-MO-001/93 (1993).
- [27] J. Santos, M. J. Rebouças and A. F. F. Teixeira, *Gen. Rel. Grav.* **27**, 989 (1995).